

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Further circulation of West Nile and Usutu virus in wild birds in Italy

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1509027> since 2016-01-27T11:58:55Z

Published version:

DOI:10.1016/j.meegid.2015.03.024

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in

Infect Genet Evol. 2015;32:292-7. doi: 10.1016/j.meegid.2015.03.024.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>),
<http://www.sciencedirect.com/science/article/pii/S1567134815001094>

Further circulation of West Nile and Usutu viruses in wild birds in Italy

Isis Victoriano Llopis ^a, Luca Rossi ^a, Annapia Di Gennaro ^c, Andrea Mosca ^b, Liana Teodori ^c, Laura Tomassone ^a, Elena Grego ^a, Federica Monaco ^c, Alessio Lorusso ^{c,*}, Giovanni Savini ^c

^a Dipartimento di Scienze Veterinarie, University of Turin, Italy

^b Istituto per le Piante da Legno e l'Ambiente, IPLA spa, Turin, Italy

^c OIE Reference Laboratory for West Nile Fever, Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise, Teramo, Italy

* corresponding author

Abstract

Usutu virus (USUV) and West Nile virus (WNV) are emerging pathogens that can cause neurological disease in humans. From March 2012 to June 2013, a sero-survey on wild birds was carried out to investigate the circulation of both viruses in Northwest (NW) Italy. Samples belonging to 47 different bird species have been collected using a volunteer based network and a wildlife rehabilitation center. Four of 297 serum samples had neutralizing antibodies against USUV ($P = 1.34\%$, IC 95% 0.36–3.4), while 10 of 233 samples tested positive for WNV ($P = 4.29\%$, IC 95% 2.07–7.75). Neutralizing antibodies for WNV were significantly more prevalent ($p < 0.001$) in trans-Saharan migrants ($P = 21\%$, IC 95% 9.55–37.3) than in resident and short-distance birds, but no migratory habit-related differences were found for USUV. Antibodies in resident bird species suggest that both viruses are circulating in NW Italy.

1. Introduction

Usutu virus (USUV) and West Nile virus (WNV) are emerging neuro-pathogenic agents that belong to the Japanese encephalitis virus antigenic complex of the family *Flaviviridae*, genus *Flavivirus*. Both viruses are maintained in the environment through a bird-mosquito life cycle (Hubálek, 2008) whereas mammals including humans are so far regarded as incidental or dead-end hosts. Migratory birds are assumed to have a key role in the amplification and circulation of these viruses (Malkinson and Banet, 2002; Weissenböck et al., 2002). WNV appeared for the first time in Italy in 1998 causing encephalitis in horses in Tuscany (Autorino et al., 2002). Ten years later, WNV reappeared in northern Italy affecting horses and humans (Calistri et al., 2010a,b; Monaco et al., 2010, 2011). In order to monitor and control WNV circulation, a serological, entomological and virological surveillance program for West Nile neuroinvasive disease has been implemented at the national level by the Italian Ministry of Health (Italian Ministry of Health, 2008). Within this framework it was possible to detect viral circulation in birds, mosquitoes and equids (Monaco et al., 2010, 2011; Calzolari et al., 2010; Savini et al., 2012, 2013) and several human cases were reported (Calistri et al., 2010a; Rizzo et al., 2009, 2012; Bagnarelli et al., 2011; Delbue et al., 2014). USUV was first reported in Austria in 2001 when a considerable die-off of Eurasian blackbirds (*Turdus merula*) was observed in and around Vienna, but a recent retrospective analysis of archived samples from dead birds in the Tuscany region (Italy) in 1996 provided evidence for an earlier introduction into Europe (Weissenböck et al., 2002, 2013). USUV has been noticed again in the last decade in some regions of northern Italy by virological and serological methods (Calzolari et al., 2013; Lelli et al., 2008;

Savini et al., 2011). In 2009, two cases of human encephalitis associated to USUV infection were reported in Emilia Romagna region (northern Italy) confirming the zoonotic potential of this virus (Cavrini et al., 2009; Pecorari et al., 2009).

By working in collaboration with organizations caring for wildlife the surveillance program may be improved by increasing the number and diversity of samples (Nemeth et al., 2007). Moreover, wildlife rehabilitation centers may greatly enhance and simplify surveillance efforts for avian-related viruses in some areas by concentrating many samples in limited space (Nemeth et al., 2007). Samples collected from free-ranging birds during ringing campaigns may also provide additional epidemiological information (Komar, 2000). In this perspective, we performed a serological investigation within wild birds collected in the Piedmont region in order to investigate the circulation of WNV and USUV viruses. Furthermore, we evaluated how the use of serological investigation from wild birds, obtained by volunteer networks, may integrate the data derived from official surveillance protocols.

2. Materials and methods

2.1. Study area and sampling sites

The study was carried out in Piedmont, a region of Northwest (NW) Italy (Fig. 1A). From March 2012 to June 2013, 304 blood samples were collected from wild birds belonging to 47 different species (Table 1A and B). Of these, 168 individuals were captured using mist-nets placed in two different ringing stations: Scrivia River Valley (N = 147) (province of Alessandria, 44.8087 N, 8.8572 E) and San Genuario Marsh reserve (N = 21) (province of Vercelli, 45.2175 N, 8.1777 E). These locations were selected based upon the high ecological richness and the abundance of mosquitoes (Pollono et al., 1998). Several bird species often breed in both locations, and, remarkably, the Scrivia River Valley is along one of the main migratory paths between Europe and Africa (Silvano and Boano, 2008). Captured birds were identified according to species, sex and age class (Spina and Volponi, 2008a). Birds were then ringed, sampled and released. Other blood samples (N = 136) were made available by the C. A. N. C. (“Centro recupero animali non convenzionali”), a wildlife rehabilitation center at the Department of Veterinary Sciences, University of Turin, which hospitalizes rescued birds from several areas of NW Italy. Samples origin is shown in Fig. 1B.

Birds were classified in one of the following three groups according to their migratory habits, as indicated in Table 1A and B: residents, short-distance migrants and trans-Saharan migrants (Spina and Volponi, 2008a,b). For those species with a mixed behavior, prevalence data were treated independently and attributed each time to one and to the other behavior group.

2.2. Sampling procedure

Blood samples were collected by syringes or capillary tubes for micro-hematocrit from the brachial or jugular veins according to species. The volume of collected blood never exceeded the 1% of body mass (McGuill and Rowan, 1989). Samples were allowed to clot at room temperature and then centrifuged for 10 min at 5600 g for Eppendorf tubes (Eppendorf Srl Milan, Italy) and 5 min at 3500 g for micro-hematocrit tubes. Sera were stored at -20°C until use.

2.3. Laboratory tests

2.3.1. Virus strains

USUV strain 939/01 isolated from a blackbird in Vienna (Austria) in 2001 and WNV strain Eg-101 were kindly donated by Prof Zdenek Hubalek (Medical Zoology Laboratory, Institute of Vertebrate Biology, Academy of Sciences, Valtice, Czech Republic) and the Unité des Arbovirus et des Fièvres hémorragiques, Institut Pasteur, Paris (France), respectively. The two viruses are routinely used for the diagnostic activities at the Istituto Zooprofilattico Sperimentale of Teramo.

2.3.2. Serological investigation

A total number of 304 serum samples were tested by serum-neutralization (SN) assay according to a recent protocol developed by our group (Di Gennaro et al., 2014). The small volume of some sera also influenced the diagnostic pipeline. In particular, out of 304 samples, 233 and 297 serum samples were tested for the presence of WNV and USUV neutralizing antibodies, respectively. Of these, 226 samples were tested simultaneously for WNV and USUV. In few cases (N = 34) the serological screening for USUV was performed starting at 1:20 dilution of the tested serum.

2.3.3. Molecular detection of WNV and USUV

Two real-time RT-PCR assays were employed for the molecular detection of WNV (Del Amo et al., 2013) and USUV (Cavrini et al., 2007). Nucleic acids were purified from the blood samples of serological positive birds by means of BioSprint 96 One-For-All Vet Kit (QIAGEN, Germany).

3. Results

Results of the serological analyses are shown in Table 2. Out of 297 samples tested for USUV, four had specific neutralizing antibodies (1.34%; 95% IC 0.36–3.4). Antibodies were detected from a blackbird (*T. merula*), a long eared owl (*Asio otus*), a rock pigeon (*Columba livia*) and from a mallard (*Anas platyrhynchos*), sampled at the municipalities of Villarvernia, Scalenghe, Grugliasco and Castagnole Piemonte, respectively (Fig. 1C). Neutralising titers ranged from 1:10 to 1:40. The prevalence of USUV sero-reactors was similar despite the migratory habits of the sampled birds. However, USUV antibodies prevalence may have been underestimated as the serological screening of some sera started at 1:20 dilution.

Out of 233 serum samples tested for WNV, ten had neutralizing antibodies (4.29%; IC 95% 2.07–7.75). The positive samples belonged to a Eurasian scops owl (*Otus scops*), two little bittern (*Ixobrychus minutus*), one nighthale (*Luscinia megarhynchos*), one Eurasian hobby (*Falco subbuteo*), one European green woodpecker (*Picus viridis*) and one common kestrel (*Falco tinnunculus*), captured in the municipalities of Vercelli, Buriasso, Villarvernia, Trofarello and Villarpellice, respectively (Fig. 1D). The neutralizing titers ranged from 1:10 to 1:80. Neutralizing antibodies for WNV were significantly more prevalent ($p < 0.001$) in trans-Saharan migrants ($P = 21\%$, IC 95% 9.55–37.3, N = 38) than in resident ($P = 1.06\%$, IC 95% 0.02–5.78, N = 94) or short-distance birds ($P = 0.99\%$, IC 95% 0.02–5.39, N = 101). Serological positive samples for WNV (10) and USUV (4) belonged to the 226 serum samples tested by SN for both viruses. Cross-

reactions have not been detected. WNV or USUV serological positive animals turned out to be negative by molecular assays for WNV and USUV RNA.

4. Discussion

In the present manuscript, the exposure to USUV and WNV viruses was investigated in several wild bird species in Piedmont region, northwestern Italy (NW). At the time the present surveillance in wild birds was ongoing (March 2012–June 2013), WNV was never demonstrated to circulate in Piedmont region, and only 3 pools collected during 2009 and 2010 of USUV-positive *Culex pipiens* were reported in a peripheral eastbound area of the region (Cerutti et al., 2012, Fig. 1C). Therefore the aim of the study was primarily to set up a surveillance network based on serological investigation of blood samples collected from live wild birds in order to explore the prevalence of WNV and USUV antibodies in that area.

The scenario slightly changed in the following years. Indeed, WNV was revealed in the region by molecular methods in pools of *C. pipiens* at the end summer of 2014 confirming therefore the circulation of WNV as suggested in the present study (Dr. Cristina Casalone, Istituto Zooprofilattico of Piemonte, Liguria and Val d'Aosta, <http://promedmail.chip.org/pipermail/promed/2014-November/005857.html>, Fig. 1D). Moreover, a recent paper (Rizzo et al., 2014) describes the evidence of USUV circulation within mosquitoes in Piedmont region (Novara province) during 2011 and 2012 (Fig. 1C). This area is in very close proximity to the Ticino river natural park, a wet area and an important wintering site for wild waterfowl. This recent data and the results produced in this study confirm the establishment of an enzootic cycle of USUV in the Piedmont region.

This survey revealed antibodies against WNV and USUV in resident bird species also. Blood samples from serological positive animals were further screened by molecular assays in order to detect active viraemia. However, none of them showed WNV or USUV RNA in the bloodstream. It was not possible to test the remaining samples as for the limited amount of material to be processed for RNA purification. Reasonably, our study shows some limitations since we were not able to demonstrate active circulation of WNV and USUV in the sampled wild birds. However, the present serological investigation combined with the detection of WNV or USUV RNA from mosquito pools described by other research groups dynamically working on flavivirus circulation in Piedmont region, strongly suggests active circulation of both viruses.

It is assumed that USUV has been introduced in Europe by trans-Saharan migratory species and that the virus adapted over the time to the local environment by establishing a transmission cycle among local birds and mosquitoes (Chvala et al., 2007). On the other hand, the transmission within European countries may have occurred by short-distance migratory birds (Steinmez et al., 2011). Alternatively, trans-Saharan migratory birds may be responsible for the introduction of USUV every year in Europe (Buckley, 2003). The results of this study are in contrast with the second assumption, as no migratory birds tested positive for USUV antibodies and all positive birds were resident or short-distance migrants. Regarding the introduction of WNV, our study supports the scenario of a mixed system in which long-distance migrant birds are primarily involved in spreading WNV from Africa to Europe whereas resident and short distance migratory birds

contribute, afterwards, to the establishment of local enzootic cycles (Savini et al., 2013; Monaco et al., 2011).

Though USUV exposed birds were scattered over a relatively wide area (Fig. 1C), the prevalence of sero-reactors suggests a limited circulation of the virus. Nevertheless, the 1.34% prevalence found is within the range reported from wild birds in Italy (0– 5.2%) (Lelli et al., 2008; Savini et al., 2011). So far, USUV outbreaks with fatal neurological outcome in wild birds have not been reported in NW Italy. In general, the antibody titers against USUV were low, ranging between 1:10 and 1:40, but in naturally and experimentally infected birds, flaviviral antibody titers are generally low and prone to considerable variations within short time (Savini et al., 2011). We may speculate on the putative locations where seropositive birds were exposed to USUV, on the basis of their migratory or sedentary behavior. Two scenarios can be indeed estimated for the juvenile common blackbird sampled in late summer: it may have been exposed while residing in the study area or, alternatively, the presence of specific USUV antibodies derived from passive transfer of maternal immunity, as previously described for WNV (Gibbs et al., 2005). The exposure site of the long eared owl is questionable, since individuals captured in NW Italy are partly breeding or transient (Fasano et al., 2005). The remaining USUV sero-positive birds were one mallard and one rock pigeon. Mallards reside in most of the European countries except in the North and East of Europe, from where they migrate during the winter season (Spina and Volponi, 2008a). Migratory individuals usually arrive in the Piedmont region from northern Europe in November lasting until February-March (Fasano et al., 2005). Possibly, the sero-positive mallard was a resident individual that became infected in NW Italy as the capture occurred in April when wintering mallards are usually gone. However, the presence of USUV antibodies in a rock pigeon, a resident species, suggests a local circulation of the virus.

Antibodies against WNV were detected in two little bitterns sampled few weeks before migration period toward wintering areas. One of them had a high antibody titer (1:80) suggesting a local recent exposure to WNV. High antibody titers (1:80) were also revealed in the nightingale and in one of the Eurasian scops owls. Their arrival in NW Italy from the wintering regions usually occurs in April-May (Fasano et al., 2005). Since both birds were sampled in May, it is reasonable to assume that they were exposed prior to migrate to Europe. The common kestrel was the only short distance migrant among the WNV sero-reactors, but in NW Italy there are also described to be as resident individuals (Fasano et al., 2005). Because of this, the exposure site of the kestrel is difficult to identify. The positive European green woodpecker, a resident species, is a further evidence of a local WNV circulation, as demonstrated elsewhere in other Italian regions (Savini et al., 2013). Significant higher WNV sero-prevalence was detected in trans-Saharan migrants, in accordance with other surveys (López et al., 2008). This difference suggests that long-distance migrants are exposed during the journeys and/or during the winter stay in Africa to a greater magnitude of WNV circulation as compared to the breeding grounds in Europe.

Overall, antibody titers against WNV and USUV detected in wild birds were similar to those regularly observed during experimental infections in birds with WNV (Del Amo et al., 2014; Pérez-Ramírez et al., 2014). It is noteworthy that a slightly lower (1:10–1:40 titer range) neutralizing antibody response was detected for USUV with respect to the neutralizing antibody response observed in those wild birds found serologically positive for WNV showing titers ranging from

1:10 to 1:80. This slight difference in antibody titers could be consistent with a less intense infection by USUV compared with WNV in wild birds.

This study offers a further evidence of the local circulation of WNV and USUV within wild bird species in the NW of Italy. We argue that sampling of live birds allowed a better representation of both viruses distribution for two reasons. First, our survey evaluated a wide range of potential USUV and WNV amplifier hosts mostly dwelling in rural and sylvatic habitats. The chance to detect an emerging infection in these habitats through passive surveillance is low compared with urban areas (Marra et al., 2004), even more so when lethality of circulating viruses is low or when herd immunity is present (Buckley, 2003). Second, surveillance national plans based on both active and passive surveillances are primarily focused on providing direct evidence of early viral circulation (Italian Ministry of Health, 2008) by introducing sentinel fowl in a given area supported by the entomological surveillance and by a very limited number of resident synanthropic wild birds mostly originating from urban areas.

Results of this survey prove that ringing organizations and wildlife rehabilitation centers may be fruitfully involved during the surveillance of bird-related emerging or re-emerging pathogens, as already previously stressed (Nemeth et al., 2007). This – mostly voluntary – network is widespread in Europe (Euring, 2013) and may be potentially very active. In NW Italy, for instance, 44 accredited ringers had worked in 116 stations during 2012. They handled in total 35,660 wild birds belonging to 151 different species (Fasano et al., 2013). In addition, one of the six regional wildlife rehabilitation centers present in the Piedmont region provided samples for the present investigation thus giving evidence of their potential role for a robust surveillance plan assessment. In a public health perspective, the contribution of this network to WNV surveillance would be of particular interest in areas which are not officially infected and/or are close to endemic areas. Data from this network would be of great value to reveal additional information upon WNV and USUV circulation and subsequently address other costly and labor-intensive actions specifically targeted to human health protection. Investigations on the occurrence, ecology and epidemiology of these two important zoonotic viruses are paramount. In this perspective, our work further highlights the need for the establishing of tight connections between veterinarians, physicians, entomologists, public health officers and dynamic voluntary networks.

Acknowledgments

The authors deeply thank Fabrizio Silvano, Piero Bravin, Sergio Fasano and ringing stations volunteers for assisting in bird captures and sample collection. We also thank Dr. Alessandro Starvaggi and Dr. Andrea Dogliero, veterinary personnel of C.A.N.C., for providing blood samples. Funding was provided by the Italian Ministry of Health. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the Istituto Zooprofilattico Sperimentale dell'Abruzzo e Molise.

References

- Autorino, G.L., Battisti, A., Deubel, V., Ferrari, G., Forletta, R., Giovannini, A., Lelli, R., Murri, S., Scicluna, M.T., 2002. West Nile virus epidemic in horses, Tuscany region, Italy. *Emerg. Infect. Dis.* 8, 1372–1378.
- Bagnarelli, P., Marinelli, K., Trotta, D., Monachetti, A., Tavio, M., Del Gobbo, R., Capobianchi, M., Menzo, S., Nicoletti, L., Magurano, F., Varaldo, P., 2011. Human case of autochthonous West Nile virus lineage 2 infection in Italy, September 2011. *Eurosurveillance* 16, pii: 20002.
- Buckley, a., 2003. Serological evidence of West Nile virus, Usutu virus and Sindbis virus infection of birds in the UK. *J. Gen. Virol.* 84, 2807–2817.
- Calistri, P., Giovannini, A., Savini, G., Monaco, F., Bonfanti, L., Ceolin, C., Terregino, C., Tamba, M., Cordioli, P., Lelli, R., 2010a. West Nile virus transmission in 2008 in north-eastern Italy. *Zoonoses Public Health* 57, 211–219.
- Calistri, P., Monaco, F., Savini, G., Guercio, A., Purpari, G., Vicari, D., Cascio, S., Lelli, R., 2010b. Further spread of West Nile virus in Italy. *Vet. Ital.* 46, 467–474.
- Calzolari, M., Bonilauri, P., Bellini, R., Albieri, A., Defilippo, F., Maioli, G., Galletti, G., Gelati, A., Barbieri, I., Tamba, M., Lelli, D., Carra, E., Cordioli, P., Angelini, P., Dottori, M., 2010. Evidence of simultaneous circulation of West Nile and Usutu viruses in mosquitoes sampled in Emilia-Romagna region (Italy) in 2009. *PLoS ONE* 5, e14324.
- Calzolari, M., Bonilauri, P., Bellini, R., Albieri, A., Defilippo, F., Tamba, M., Tassinari, M., Gelati, A., Cordioli, P., Angelini, P., Dottori, M., 2013. Usutu virus persistence and West Nile virus inactivity in the Emilia-Romagna region (Italy) in 2011. *PLoS ONE* 8, e63978.
- Cavrini, F., Della Pepa, M.E., Gaibani, P., Pierro, A.M., Rossini, G., Landini, M.P., Sambri, V., 2007. A rapid and specific real-time RT-PCR assay to identify Usutu virus in human plasma, serum, and cerebrospinal fluid. *J. Clin. Virol.* 50, 221–223.
- Cavrini, F., Gaibani, P., Longo, G., Pierro, A.M., Rossini, G., Bonilauri, P., Gerunda, G.E., Gerundi, G.E., Di Benedetto, F., Pasetto, A., Girardis, M., Dottori, M., Landini, M.P., Sambri, V., 2009. Usutu virus infection in a patient who underwent orthotopic liver transplantation, Italy, August–September 2009. *Euro Surveill.* 14, 17–18.
- Cerutti, F., Giacobini, M., Mosca, A., Grasso, I., Rambozzi, L., Rossi, L., Bertolotti, L., 2012. Evidence of mosquito-transmitted flavivirus circulation in Piedmont, north-western Italy. *Parasit. Vectors* 5, 99.
- Chvala, S., Bakonyi, T., Bukovsky, C., Meister, T., Brugger, K., Rubel, F., Nowotny, N., Weissenböck, H., 2007. Monitoring of Usutu virus activity and spread by using dead bird surveillance in Austria, 2003–2005. *Vet. Microbiol.* 122, 237–245.
- Del Amo, J., Sotelo, E., Fernández-Pinero, J., Gallardo, C., Llorente, F., Agüero, M., Jiménez-Clavero, M.A., 2013. A novel quantitative multiplex real-time RT-PCR for the simultaneous

detection and differentiation of West Nile virus lineages 1 and 2, and of Usutu virus. J. Virol. Methods 189, 321–327.

Del Amo, J., Llorente, F., Pérez-Ramírez, E., Soriguer, R.C., Figuerola, J., Nowotny, N., Jiménez-Clavero, M.A., 2014. Experimental infection of house sparrows (*Passer domesticus*) with West Nile virus strains of lineages 1 and 2. Vet. Microbiol. 172, 542–547.

Delbue, S., Ferrante, P., Mariotto, S., Zanusso, G., Pavone, A., Chinaglia, M., L'Erario, R., Monaco, S., Ferrari, S., 2014. Review of West Nile virus epidemiology in Italy and report of a case of West Nile virus encephalitis. J. Neurovirol. 20, 437–441.

Di Gennaro, A., Lorusso, A., Casaccia, C., Conte, A., Monaco, F., Savini, G., 2014. Serum-neutralization assay can efficiently replace plaque reduction neutralization test for detection and quantitation of West Nile Virus antibodies in human and animal serum samples. Clin. Vaccine Immunol. 21, 1460–1462.

Euring, 2013. Co-ordinating bird ringing throughout Europe. <<http://www.euring.org>>.

Fasano, S., Boano, G., Ferro G., 2005. 25 anni di inanellamento in Piemonte e Valle d'Aosta. Museo Civico Craveri di Storia Naturale. Memorie dell'ANP, vol. V. Bra. pp. 224.

Fasano, S., Mervic, C., Ranghino, S., 2013. Resoconto dell'attività di inanellamento degli uccelli a scopo scientifico in Piemonte e Valle d'Aosta Anno 2012. Riv. Piemont. Stor. Nat. 34, 367–387.

Gibbs, S.E.J., Hoffman, D.M., Stark, L.M., Marlenee, N.L., Blitvich, B.J., Beaty, B.J., Stallknecht, D.E., 2005. Persistence of antibodies to West Nile Virus in naturally infected rock pigeons. Clin. Vaccine Immunol. 12, 665–667.

Hubálek, Z., 2008. Mosquito-borne viruses in Europe. Parasitol. Res. 103, 29–43. Italian Ministry of Health. Approval of the national surveillance plan for West Nile encephalomyelitis [in Italian]. Rome: Ministry of Health, 12 February 2008. G. U. Serie Generale no. 36. <<http://www.normativasanitaria.it/jsp/dettaglio.jsp?aggiornamenti=&attoCompleto=si&id=25243&page=&anno=null>>.

Komar, N., 2000. West Nile virus surveillance using sentinel birds. Ann. N. Y. Acad. Sci. 951, 58–73.

Lelli, R., Savini, G., Teodori, L., Filipponi, G., Di Gennaro, a., Leone, a., Di Gialleonardo, L., Venturi, L., Caporale, V., 2008. Serological evidence of USUTU virus occurrence in north-eastern Italy. Zoonoses Public Health 55, 361–367.

López, G., Jiménez-Clavero, M.A., Tejedor, C.G., Soriguer, R., Figuerola, J., 2008. Prevalence of West Nile virus neutralizing antibodies in Spain is related to the behavior of migratory birds. Vector Borne Zoonotic Dis. 8, 615–621.

Malkinson, M., Banet, C., 2002. The role of birds in the ecology of West Nile virus in Europe and Africa. Curr. Topic Microbiol. Immunol. 267, 309–322.

- Marra, P.P., Griffing, S., Caffrey, C., Kilpatrick, a.M., McLean, R., Brand, C., Saito, E., Dupuis, A.P., Kramer, L., Novak, R., 2004. West Nile virus and wildlife. *Bioscience* 54, 393.
- McGuill, M.W., Rowan, A.N., 1989. Biological effects of blood loss: implications for sampling volumes and techniques. *ILAR J.* 31, 5–20.
- Monaco, F., Lelli, R., Teodori, L., Pinoni, C., Di Gennaro, A., Polci, A., Calistri, P., Savini, G., 2010. Re-emergence of West Nile virus in Italy. *Zoonoses Public Health* 57, 476–486.
- Monaco, F., Savini, G., Calistri, P., Polci, A., Pinoni, C., Bruno, R., Lelli, R., 2011. 2009 West Nile disease epidemic in Italy: first evidence of overwintering in Western Europe? *Res. Vet. Sci.* 91, 321–326.
- Nemeth, N.M., Beckett, S., Edwards, E., Klenk, K., Komar, N., 2007. Avian mortality surveillance for West Nile virus in Colorado. *Am. J. Trop. Med. Hyg.* 76, 431–437.
- Pecorari, M., Longo, G., Gennari, W., Grottola, A., Sabbatini, A.M.T., Tagliazucchi, S., 2009. Rapid communications: First human case of Usutu virus neuroinvasive infection, Italy, August–September 2009. *Euro Surveill.* 14, 15–16.
- Pérez-Ramírez, E., Llorente, F., Jiménez-Clavero, M.Á., 2014. Experimental infections of wild birds with West Nile virus. *Viruses* 6, 752–781.
- Pollono, F., Rossi, L., Cancrini, G., 1998. Indagine sui Culicidi da esca canina in Piemonte. *Parassitologia* 40, 439–445.
- Rizzo, C., Vescio, F., Declich, S., Finarelli, A.C., Macini, P., Mattivi, A., Rossini, G., Piovesan, C., Barzon, L., Palù, G., Gobbi, F., Macchi, L., Pavan, A., Magurano, F., Ciufolini, Nicoletti, L., Salmaso, S., Rezza, G., 2009. West Nile virus transmission with human cases in Italy, August–September 2009. *Euro Surveill.* 14, 5–8.
- Rizzo, C., Salcuni, P., Nicoletti, L., Ciufolini, M.G., Russo, F., Masala, R., Frongia, O., Finarelli, A.C., Gramegna, M., Gallo, L., Pompa, M.G., Rezza, G., Salmaso, S., Declich, S., 2012. Epidemiological surveillance of west Nile neuroinvasive diseases in Italy, 2008 to 2011. *Eurosurveillance* 17, pii: 20172.
- Rizzo, F., Cerutti, F., Ballardini, M., Mosca, A., Vitale, N., Radaelli, M.C., Desiato, R., Prearo, M., Pautasso, A., Casalone, C., Acutis, P., Peletto, S., Mandola, M.L., 2014. Molecular characterization of flaviviruses from field-collected mosquitoes in northwestern Italy, 2011–2012. *Parasit. Vectors* 7, 395.
- Savini, G., Monaco, F., Terregino, C., Di Gennaro, A., Bano, L., Pinoni, C., De Nardi, R., Bonilauri, P., Pecorari, M., Di Gialleonardo, L., Bonfanti, L., Polci, A., Calistri, P., Lelli, R., 2011. Usutu virus in Italy: an emergence or a silent infection? *Vet. Microbiol.* 151, 264–274.
- Savini, G., Capelli, G., Monaco, F., Polci, A., Russo, F., Di Gennaro, A., Marini, V., Teodori, L., Montarsi, F., Pinoni, C., Pisciella, M., Terregino, C., Marangon, S., Capua, I., Lelli, R., 2012. Evidence of West Nile virus lineage 2 circulation in Northern Italy. *Vet. Microbiol.* 158, 267–273.

Savini, G., Puggioni, G., DI Gennaro, A., DI Francesco, G., Rocchigiani, A.M., Polci, A., Marini, V., Pinoni, C., Rolesu, S., Marruchella, G., Lorusso, A., Monaco, F., 2013. West Nile virus lineage 2 in Sardinian wild birds in 2012: a further threat to public health. *Epidemiol. Infect.* 141, 2313–2316.

Silvano, F., Boano, G., 2008. Gli uccelli della Valle Scrivia e il centro di inanellamento della Provincia di Alessandria, Ed. Litografia Viscardi (AL). pp. 15–20.


Spina, F., Volponi, S., n.d. Atlante della Migrazione degli Uccelli in Italia II. Passeriformi. Spina, F., Volponi, S., 2008(b). Atlante della Migrazione degli Uccelli in Italia. 1. non-Passeriformi. Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA). Tipografia CSR-Roma. 800 pp.

Spina, F., Volponi, S., 2008(a). Atlante della Migrazione degli Uccelli in Italia. 2. Passeriformi. Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA). Tipografia SCR-Roma. 632 pp.

Steinmez, H.W., Bakonyi, T., Weissenböck, H., Hatt, J.M., Eulenberger, U., Robert, N., Hoop, R.K., Nowotny, N., 2011. Emergence and establishment of Usutu virus infection in wild and captive avian species in and around Zurich, Switzerland - genomic and pathologic comparison to other central European outbreaks. *Vet. Microbiol.* 148, 207–212.

Weissenböck, H., Kolodziejek, J., Url, A., Lussy, H., Rebel-bauder, B., Nowotny, N., 2002. Emergence of USUTU virus, an African Mosquito-Borne Flavivirus of the Japanese Encephalitis Virus Group, Central Europe. *Emerg. Infect. Dis.* 8, 652–656.

Weissenböck, H., Bakonyi, T., Rossi, G., Mani, P., Nowotny, N., 2013. Usutu virus, Italy, 1996. *Emerg. Infect. Dis.* 19, 274–277

Fig. 1. Map of Italy highlighting the geographical position of Piedmont region (A). Map of Piedmont region showing the geographical origin of the samples (B). Geographical origin of USUV positive samples: municipalities of Villarvernia (VV), Scalenghe (S), Grugliasco (G) and Castagnole Piemonte (CP) (C) and positive samples for WNV: municipalities of Vercelli (VE), Buriasco (B), Villarvernia (VV), Trofarello (T) and Villarpellice (VP) (D).  : Geographical location of mosquitoes' pools positive for USUV (B) and WNV (C) RNA demonstrated in 2009/2010, 2012 and 2014.




Table 1. (A) and (B) Species, number of sampled individuals tested for USUV and WNV. Each species was classified according to their migratory behavior in resident (R), short-distance migrant (M) or trans-Saharan migrant (T). Birds were divided in two tables, (A) and (B), according to the alphabetical order.

Table 1A

	Migratory behavior	Total samples	Number of samples tested for WNV	Number of samples tested for USUV
<i>Accipiter gentilis</i>	R	2	1	2
<i>Accipiter nisus</i>	M	1	1	1
<i>Alcedo atthis</i>	R - M	1	0	1
<i>Anas platyrhynchos</i>	R - M	6	4	6
<i>Anser anser</i>	M	1	1	1
<i>Apus apus</i>	T	1	1	1
<i>Ardea cinerea</i>	M	3	3	3
<i>Asio otus</i>	M	5	4	5
<i>Athene noctua</i>	R	3	2	3
<i>Botaurus stellaris</i>	M	1	1	1
<i>Bubo bubo</i>	R	1	1	1
<i>Buteo buteo</i>	M	7	7	7
<i>Cairina moschata</i>	R	2	2	2
<i>Caprimulgus europaeus</i>	M	3	2	2
<i>Chroicocephalus ridibundus</i>	M	4	4	4
<i>Columba livia</i>	R	45	41	45
<i>Columba palumbus</i>	M	4	4	4
<i>Corvus cornix</i>	R	23	22	23
<i>Corvus monedula</i>	M	1	1	1
<i>Coturnix coturnix</i>	T	1	1	1
<i>Cygnus olor</i>	M	1	1	1
<i>Egretta garzetta</i>	R - M	2	1	2
<i>Erithacus rubecula</i>	M	1	0	1
<i>Falco subbuteo</i>	T	1	1	1
<i>Falco tinnunculus</i>	R - M	6	3	6

Table 1B

	Migratory behavior	Total samples	Number of samples tested for WNV	Number of samples tested for USUV
<i>Garrulus glandarius</i>	R - M	7	7	7
<i>Ixobrychus minutus</i>	T	6	3	6
<i>Lanius collurio</i>	T	2	2	2
<i>Larus michahellis</i>	R - M	2	1	2
<i>Luscinia megarhynchos</i>	T	3	1	3
<i>Merops apiaster</i>	T	20	10	18
<i>Otus scops</i>	T	18	10	18
<i>Parus major</i>	R - M	6	3	5
<i>Passer italiae</i>	R	1	1	1
<i>Phoenicurus phoenicurus</i>	T	3	1	3
<i>Picoides major</i>	R - M	5	3	5
<i>Pica pica</i>	R	15	14	15
<i>Picus viridis</i>	R	4	3	4
<i>Pyrrhocorax pyrrhocorax</i>	R	1	1	1
<i>Strix aluco</i>	R	3	3	3
<i>Streptopelia decaocto</i>	R	4	3	4
<i>Streptopelia turtur</i>	T	8	7	8
<i>Sturnus vulgaris</i>	M	38	32	38
<i>Sylvia atricapilla</i>	R - T	2	0	2
<i>Tachymarptis melba</i>	T	1	1	1
<i>Turdus merula</i>	M	28	18	25
<i>Turdus philomelos</i>	M	1	0	1
Total number		304	233	297

Table 2

Titers and migratory behavior for individual birds with USUV and WNV neutralizing antibodies sampled between March 2012 and June 2013 in Piedmont, Italy. Each species was classified as resident (R), short-distance migrant (M), or trans-Saharan migrant (T).

[illegible]